

PARALLEL PROJECTION AREA AND HOLDITCH'S THEOREM

H. BAYRAM KARADAĞ and SADIK KELEŞ

Department of Mathematics, Faculty of Science and Arts, İnönü University, Malatya, Turkey

(Received May 24, 1996; Accepted Nov. 4, 1996)

ABSTRACT

In this paper, the parallel projection area of a closed spatial curve formed under the motion $B(c_1)$ defined along the closed spherical curve c_1 [12] have been calculated. After that Holditch's Theorem [7] and its some corollaries which is well-known [3] have been generalized to closed spatial curve.

1. INTRODUCTION

The study of one-parameter closed motions become an interesting subject in kinematics after the work of Jacob Steiner [11] and H. Holditch [7].

During the second half of the nineteenth century, there appeared many publications about Steiner's and Holditch's Theorems; for example: C. Leudesdorf [9, 10] and A.B. Kempe [8].

After the work of Steiner and Holditch the first study about spherical motions was given by E.B. Elliott [2, 3, 4]. Another study in this field was also given by H.R. Müller [11]. H.H. Hacısalihoğlu [6] obtained a formula which is equivalent to the Holditch formula. R. Güneş and S. Keleş [5], using the area formula and the area vector given by W. Blaschke [1] and H.R. Müller [11], respectively, obtained the formula given by H.H. Hacısalihoğlu by a different method.

H. Pottmann [12] defined the spherical motion along a curve on a sphere and also he gave the parallel projection area of the spherical indicator using the parallel projection area vector.

In this study, H. Holditch's Theorem, which is well-known for one-parameter closed planar motions, was generalized to the closed spatial motions by using the area vector which was described by H.R. Müller

[11] and parallel projection area formula which was given by H. Pottmann [12] and some results were obtained.

2. SPHERICAL CURVES

Let a c_1 -curve of class C^2 on a unit sphere K^1 of 3-dimensional Euclidean space is given by

$$\begin{aligned} \vec{c}_1: t \in I \subset \mathbb{R} \rightarrow \vec{c}_1(t) \in \mathbb{R}^3, \|\vec{c}_1\| = 1, \vec{c}_1 \in C^2(I) \\ c_1 = \vec{c}_1(I) \end{aligned} \quad (1)$$

where C^2 denotes the set of twice continuously differentiable curves.

Let us consider the sphere K coinciding with K^1 to be $K = K^1$, where K^1 is a fixed sphere and K is a moving sphere with respect to K^1 . In this case the curve c_1 on K^1 defines an accompanying motion $B(c_1)$. The end point $E_1(t) \in K$ of the vector $\vec{c}_1(t)$ lies on the curve c_1 which is always tangent any constant big circles at $E_1(t)$ as illustrated in Figure 1.




Fig. 1

At the initial time, given an orthonormal frame $\{0; \vec{c}_1(t_0), \vec{c}_2(t_0), \vec{c}_3(t_0)\}$, which are rightly linked to the origin point of the moving sphere K , is defined as

$$\vec{c}_2(t_0) = \frac{\dot{\vec{c}}_1(t_0)}{\|\dot{\vec{c}}_1(t_0)\|}, \vec{c}_3(t_0) = \vec{c}_1(t_0) \wedge \vec{c}_2(t_0) \quad (2)$$

During the closed motion $B(c_1)$, a frame $\{\vec{e}_1, \vec{e}_2, \vec{e}_3\}$ can be defined for the point $E_1(t)$ of the curve c_1 at $t=t_1$ with the help of the limit $t \rightarrow t_1$. The closed motion $B(c_1)$ defined along the curve c_1 having completely the inflection point $2n$ ($n \in \mathbb{N} \cup \{0\}$) of c_1 is known as a closed motion [12].

Derivative equations of the moving frame can be written in the matrix form as

$$\begin{bmatrix} \dot{\vec{e}}_1 \\ \dot{\vec{e}}_2 \\ \dot{\vec{e}}_3 \end{bmatrix} = \begin{bmatrix} 0 & \lambda & 0 \\ -\lambda & 0 & \mu \\ 0 & -\mu & 0 \end{bmatrix} \begin{bmatrix} \vec{e}_1 \\ \vec{e}_2 \\ \vec{e}_3 \end{bmatrix} \quad (3)$$

In equation (3), Darboux rotation vector is

$$\vec{w} = \mu \vec{e}_1 + \lambda \vec{e}_3 \quad (4)$$

Hence equation (3) can be also written in the following vector form

$$\dot{\vec{e}}_i = \vec{w} \wedge \vec{e}_i, \quad (i=1,2,3) \quad (5)$$

If $\vec{w} \neq 0$ then we write the vectors

$$\vec{p}_1 = \frac{\vec{w}}{\|\vec{w}\|}, \quad \vec{p}_2 = -\frac{\vec{w}}{\|\vec{w}\|}.$$

The vectors \vec{p}_1 and \vec{p}_2 define two constant Pole points M_i , $i=1,2$, which are symmetric with respect to the origin point O , of the closed motion $B(c_1)$ at the time t . The points M_1 and M_2 are on the big circle q in the plane (\vec{e}_1, \vec{e}_3) .

In the case of $\vec{w} = \vec{0}$, an instantaneous standstill of the closed motion $B(c_1)$ is happened. During the closed motion $B(c_1)$, q is rolled without sliding on spherical evaluate q^1 of the curve c_1 .

Now, we discuss the close curve in \mathbb{R}^3 by the closed motion $B(c_1)$. Because, there is a very close relationship between spherical accompanying curves and curves theory in \mathbb{R}^3 .

Let us now consider the closed accompanying motion $B(c_1)$ that leaves origin point of a moving space R with respect to a fixed space R^1 unchanged. In such a way that we again have the derivative equations given by (3) for a space curve k in \mathbb{R}^3 .

The end points of vectors \vec{e}_1 , \vec{e}_2 and \vec{e}_3 draw, respectively, spherical tangent indicator c_1 , principal normal indicator c_2 and binormal indicator c_3 of the space curve $k \subset \mathbb{R}^3$ of the class C^1 is given as

$$\begin{aligned} \vec{x}: t \in I = [0; L) \subset \mathbb{R} \rightarrow \vec{x}(t) \in \mathbb{R}^3, \vec{x} \in C^1(\mathbb{R}) \\ \vec{x}(t+L) = \vec{x}(t) \\ k = \vec{x}(I) \end{aligned} \quad (7)$$

3. PARALLEL PROJECTION AREA AND HOLDITCH'S THEOREM

Definition 3.1. Let $c(X)$ be a closed curve in 3-dimensional Euclidean space and X be a point on $c(X)$. The vector satisfying

$$\vec{V}_X = \oint \vec{x}(t) \wedge \dot{\vec{x}}(t) dt \quad (8)$$

is called the area vector of the curve $c(X)$ [11] in which \vec{x} is the position vector of X .

Theorem 3.2. Let $c(X)$ be a closed curve in 3-dimensional Euclidean space and X be a point on $c(X)$. The projection area [11] of the planar region occurred by taking orthogonal projection onto a plane in the direction the unit vector \vec{n} of $c(X)$ is

$$2F_{X^n} = \langle \vec{n}, \vec{V}_X \rangle. \quad (9)$$

Theorem 3.3. (Holditch's Theorem [7]). If a chord of a closed curve, of constant length $a+b$, be divided into two parts of lengths a , b , respectively the difference between the areas of the closed curve, and of the locus of the dividing point, will be

$$F = F_\Gamma - F_X = \pi ab$$




Fig. 2

After the above preparations, we can calculate the area vector \vec{V}_x of the orbit $c(X)$, on \mathbb{R}^1 , of the fixed point $X \in \mathbb{R}$, during the closed accompanying motion $B(c_1)$, along a closed spherical curve c_1 . Let $I = [0, L]$ be the period interval of an instantaneous closed accompanying motion $B(c_1)$.

The position vector \vec{x} of a point $X \in \mathbb{R}$ with respect to the vectors \vec{e}_1 , \vec{e}_2 and \vec{e}_3 of K can be written as

$$\vec{x}(t) = x_1 \vec{e}_1(t) + x_2 \vec{e}_2(t) + x_3 \vec{e}_3(t) \tag{10}$$

where x_1 , x_2 and x_3 are constant coordinates of X . From (8), the area vector of the orbit $c(X)$ drawn by a point $X \in \mathbb{R}$ under the closed accompanying motion $B(c_1)$ is obtained as

$$\vec{V}_x = \sum_{i=1}^3 x_i \vec{V}_{E_i} + 2 \sum_{\substack{i,k=1 \\ i < k}}^3 x_i x_k \vec{V}_{E_{ik}} \tag{11}$$

where

$$\vec{V}_{E_i} = \int_0^L \dot{\vec{c}}_i(t) \wedge \dot{\vec{c}}_i(t) dt, \quad \vec{V}_{E_{ik}} = \frac{1}{2} \int_0^L (\dot{\vec{c}}_i(t) \wedge \dot{\vec{c}}_k(t) + \dot{\vec{c}}_k(t) \wedge \dot{\vec{c}}_i(t)) dt \tag{12}$$

We have

$$\int_0^L \dot{\vec{c}}_i(t) dt = 0 \tag{13}$$

since $B(c_1)$ is a closed accompanying motion. By using equation (3), (12) and (13) we obtain

$$\vec{V}_{E_1} = \int_0^L \lambda \dot{\vec{c}}_3(t) dt, \quad \vec{V}_{E_3} = \int_0^L \mu \dot{\vec{c}}_1(t) dt, \quad \vec{V}_{E_2} = \vec{V}_{E_1} + \vec{V}_{E_3}$$

and

$$\vec{V}_{E_{12}} = 0, \quad \vec{V}_{E_{23}} = 0, \quad \vec{V}_{E_{13}} = -\int_0^L \lambda \dot{\vec{c}}_1(t) dt = -\int_0^L \mu \dot{\vec{c}}_3(t) dt \tag{14}$$

Substitution of (14) into (11) gives

$$\vec{V}_X = \sum_{i=1}^3 x_i \vec{V}_{E_i} + 2x_1 x_3 \vec{V}_{E_{13}} \tag{15}$$

where $\vec{V}_{E_{13}}$ is the mixed area vector of the curve c_1 and the spherical curve c_3 being a spherical distance $\frac{\pi}{2}$ to c_1 .

The mixed area of orthogonal projections of curves c_1 and c_3 in the direction \vec{n} ($\|\vec{n}\| = 1$) is

$$2F_{E_{13}}^n = \langle \vec{n}, \vec{V}_{E_{13}} \rangle \quad (16)$$

Corollary 3.4. Let F_{X^n} be the projection area of the planar region occurred by taking orthogonal projection onto a plane of $c(X)$ and F_{X^p} be the projection area of the planar region happened by projecting onto same plane of $c(X)$ in any direction. From here

$$F_{X^n} = \cos \theta F_{X^p} \quad (17)$$

where θ is the angle between two image planes.

Theorem 3.5. Let the closed accompanying motion $B(c_1)$, along the curve c_1 on a unit sphere of the class C^2 , be given. Then, the orientated area F_{X^p} of parallel projection of orbit $c(X)$ of a constant point $X \in R$, in terms of the orientated areas $F_{E_{13}}^p$ and $F_{E_i}^p$, $i=1,2,3$ can be obtained as

$$F_{X^p} = 2x_1 x_3 F_{E_{13}}^p + \sum_{i=1}^3 x_i^2 F_{E_i}^p \quad (18)$$

Theorem 3.6. The orientated projection area F_{X^p} of the planar region occurred by parallel projection of the curve $c(X)$ drawn by a constant point $X \in R$ during the closed accompanying motion $B(c_1)$ is a quadratic form according to coordinates x_i , $i=1,2,3$.

If the coordinate systems are chosen properly that is, if a proper rotation is applied, from eq. (18) the orientated projection area F_{X^p} is obtained as

$$F_{X^p} = \sum_{i=1}^3 x_i^2 F_{E_i}^p \quad (19)$$

Let X and Y be two different fixed points in the moving space R . Suppose that Z is a point with the components

$$z_i = \lambda x_i + \mu y_i, \quad \lambda + \mu = 1, \quad 1 \leq i \leq 3, \quad (20)$$

on the straight line XY . The point Z has an orbit $c(Z)$ in R^1 during the closed accompanying motion $B(c_1)$. The area bounded by the orthogonal

projection of the closed curve $c(Z)$ on the plane is

$$F_{Z^p} = \lambda^2 \sum_{i=1}^3 F_{E_1^p} x_i^2 + 2\lambda\mu \sum_{i=1}^3 F_{E_1^p} x_i y_i + \mu^2 \sum_{i=1}^3 F_{E_1^p} y_i^2 \quad (21)$$

From Eq.(12), mixed area vector of curves $c(X)$ and $c(Y)$ can be obtained as

$$\vec{V}_{XY} = \sum_{i=1}^3 x_i y_i \vec{V}_{E_1} + (x_1 y_3 + y_1 x_3) \vec{V}_{E_{13}} \quad (22)$$

The orientated mixed area $F_{X^p Y^p}$ of the planar region occurred by taking parallel projection of curves $c(X)$ and $c(Y)$ onto a plane is

$$F_{X^p Y^p} = \sum_{i=1}^3 x_i y_i F_{E_1^p} + (x_1 y_3 + y_1 x_3) F_{E_{13}^p} \quad (23)$$

By using Eq.(23) in Eq.(21) we obtain

$$F_{Z^p} = \lambda^2 \sum_{i=1}^3 x_i^2 F_{E_1^p} + 2\lambda\mu \left\{ F_{X^p Y^p} - (x_1 y_3 + y_1 x_3) F_{E_{13}^p} \right\} + \mu^2 \sum_{i=1}^3 x_i^2 F_{E_1^p} \quad (24)$$

Since

$$\sum_{i=1}^3 F_{E_1^p} (x_i - y_i)^2 = F_{X^p} - 2 \left\{ F_{X^p Y^p} - (x_1 y_3 + y_1 x_3) F_{E_{13}^p} \right\} + F_{Y^p}$$

and

$$\lambda + \mu = 1 \quad , \quad \lambda^2 = 1 - \lambda\mu \quad , \quad \mu^2 = \mu - \lambda\mu \quad (25)$$

from Eq.(24), with some manipulations, we can see that

$$F_{Z^p} = \lambda F_{X^p} + \mu F_{Y^p} - \lambda\mu \sum_{i=1}^3 F_{E_1^p} (x_i - y_i)^2 \quad (26)$$

The distance between the points X and Y can be given by the metric

$$D^2(X,Y) = \varepsilon \sum_{i=1}^3 F_{E_1^p} (x_i - y_i)^2 \quad (27)$$

such that $\varepsilon = \pm 1$.

For distinct points X , Y and Z lying on the same straight line we can write

$$D(X,Y) = D(X,Z) + D(Z,Y) \quad (28)$$

Eq.(28) can be re-written as

$$\frac{D(X,Z)}{D(X,Y)} + \frac{D(Z,Y)}{D(X,Y)} = 1$$

Since $\lambda + \mu = 1$, we can take λ and μ as

$$\lambda = \frac{D(X,Z)}{D(X,Y)}, \quad \mu = \frac{D(Z,Y)}{D(X,Y)} \quad (29)$$

Eq.(26) can be written as

$$F_{Z^p} = \frac{1}{D(X,Y)} \varepsilon \left\{ F_{X^p} D(X,Z) + F_{Y^p} D(Z,Y) \right\} - \varepsilon D(X,Z) D(Z,Y) \quad (30)$$

Suppose that the fixed point X and Y in the moving space draw same closed curve (Γ) during the closed accompanying motion $B(c_1)$. In this case, $\vec{V}_X = \vec{V}_Y$ and that's why $F_{X^p} = F_{Y^p}$. Thus the Eq.(30) reduces to

$$F_{X^p} - F_{Z^p} = \varepsilon D(X,Z) D(Z,Y) \quad (31)$$

which gives generalized Holditch's Theorem.

Let ℓ be a fixed straight line in the moving space R and let four arbitrary fixed points M, X, Y and N be on the line ℓ . During the closed accompanying motion $B(c_1)$, while the points M and N move on the same curve (Γ), the points X and Y draw the different curves $c(X)$ and $c(Y)$.

Corollary 3.7. Let F and F^1 be the areas between the parallel projections of the curves (Γ) and $c(X)$ and of the curves (Γ) and $c(Y)$, respectively. Then the ratio F/F^1 depends only on the relative positions of these four points.

Proof: According to (31), the area F^1 between the projection of the curves (Γ) and $c(Y)$ is

$$F^1 = F_{M^p} - F_{Y^p} = \varepsilon D(M,Y) D(Y,N)$$

and the area F between the projection of the curves (Γ) and $c(X)$ is

$$F = F_{M^p} - F_{X^p} = \varepsilon D(M,X) D(X,N)$$

Then, joining the last two equalities the ratio F/F^1 can be obtained as

$$\frac{F}{F^1} = \frac{D(M,X) D(X,N)}{D(M,Y) D(Y,N)} \quad \text{or} \quad \frac{F}{F^1} = \left(\frac{D(M,X)}{D(M,Y)} \right)^2 \frac{D(M,Y) D(X,N)}{D(M,X) D(Y,N)} \quad (32)$$

The invariant (32) does not depend on the curve (Γ) and length of MN. It depends only on the choice of the points X and Y on MN. Since $X \neq Y$, it follows that $\frac{D(M,Y)}{D(M,X)} \neq 1$. Denote $\beta = \frac{D(M,Y)D(X,N)}{D(M,X)D(Y,N)}$. β is the cross ratio of the four points M, X, Y and N, i.e. $\beta = (MXYN)$.

Corollary 3.7 is the re-stated form of the corollary which has been given for one-parameter closed planar motions. Thus the corollary in [6] is generalized to the points of space and spatial motions.

Theorem 3.8. Let M, N, A and B be four different fixed points in the moving space R. Suppose that the line segments MN and AB meet at the point X. Then the pairs of the points M, N and A, B are on the same curve or the areas bounded by the parallel projection of the closed orbits of the pairs on the plane P are equal if and only if

$$D(M,X)D(X,N) = D(A,X)D(X,B).$$

Corollary 3.9. Let M, N, A and B be four different fixed points in the moving space R. Suppose that the line segments MN and AB meet at the point X. If the points M, N, A and B are on the same curve (Γ). Then it is $F_{M^P} = F_{A^P}$.

REFERENCES

- [1] BLASCHKE, W., Zur Bewegungen Geometrie auf der Kugel, S.B. Heidelberger Acad. Wiss Math. Kl, No.2, (1948).
- [2] ELLIOTT, E.B., A Theorem in Areas Including Holditch's, With its Analogue in Three Dimensions, Messenger Math. 7 (1878), 150-156.
- [3] ELLIOTT, E.B., Some Theorems of Kinematics on a Sphere, Proc. London Math. Soc. 1 (1881), 47-57.
- [4] ELLIOTT, E.B., Extension to Curved Surfaces of a Former Theorem on plane areas, Messenger Math. 10 (1881), 156-158.
- [5] GÜNEŞ, R., KELEŞ, S., Closed Spherical Motions and Holditch's Theorem, Mechanism and Machine Theory, Vol. 29, No. 5 (1994), pp. 755-758.
- [6] HACISALİHOĞLU, H.H., On Closed Spherical Motions, Quarterly of Appl. Math. (1971), 269-276.
- [7] HOLDITCH, H., Geometrical Theorem, The Quarterly Journal of Pure and Applied Math. 2 (1858), 38.

- [8] KEMPE, A.B., Note on Mr. Luedesdorf's Theorem in Kinematics, Messenger Math. 7 (1878), 165-167.
- [9] LUEDES DORF, C., Theorem in Kinematics, Messenger Math. 7 (1877), 125-127.
- [10] LEUDES DORF, C., Note on the Theorem in Kinematics, Messenger Math. 8 (1878), 11-12.
- [11] MÜLLER, H.R., Erweiterung des Satzes von Holditch für Geschlossene Raunkurven, Abhandl. Braunsch. Wiss. Ges. 31 (1980), 129-135.
- [12] POTTMANN, H., Über Geschlossene Spharische Kurven, El. Math. Vol. 42 (1987), 62-68.
- [13] STEINER, J., Gesamelte Werke, Berlin (1881/1882).